

Cocker Spaniel (American, English)



LABGENVET

[MyBreedData \(Donner, 2018\)](#)

Cocker Spaniel (American)

[\(AKC, 30th in popularity, 2021\)](#)

[PRA-PRCD](#)

(21 tests)

N/N	17	81.0%
M/N	3	14.3%
M/M	1	4.7%

Progressive Rod-Cone Degeneration (PRCD)

(71 tests)

N/N	51	71.8%
M/N	19	26.8%
M/M	1	1.4%

[Degenerative Myelopathy](#)

(10 tests)

N/N	9	90.0%
M/N	1	10.0%
M/M	0	0.0%

[Exercise Induced Collapse](#)

(77 tests)

N/N	72	93.5%
M/N	5	6.5%
M/M	0	0.0%

[Familial Nephropathy](#)

(8 tests)

N/N	8	100.0%
M/N	0	0.0%
M/M	0	0.0%

[Skeletal Dysplasia 2 \(SD2\)](#)

(77 tests)

N/N	75	97.4%
M/N	2	2.6%
M/M	0	0.0%

[Von Willebrand's Disease Type2 \(vWBD2\)](#)

(71 tests)

N/N	68	95.8%
M/N	3	4.2%
M/M	0	0.0%



[MyBreedData \(Donner, 2018\)](#)

English Cocker Spaniel

[\(AKC, 52nd in popularity, 2021\)](#)

Progressive Rod-Cone Degeneration (PRCD)
(82 tests)

N/N	45	54.9%
M/N	23	28.0%
M/M	14	17.1%

Familial Nephropathy (FN)
(English Cocker Spaniel) (137 tests)

N/N	134	97.8%
M/N	3	2.2%
M/M	0	0.0%

[Cone-Rod Dystrophy \(Cord1-PRA/crd4\)](#)

(137 tests)

N/N	136	99.3%
M/N	1	0.7%
M/M	0	0.0%