

Weimaraner

(AKC, 36th in popularity, 2021)



LABGENVET

[Hyperuricosuria](#)

(10 tests)

| | | |
|-----|---|-------|
| N/N | 8 | 80.0% |
| M/N | 1 | 10.0% |
| M/M | 1 | 10.0% |

Hypomyelination

(9 tests)

| | | |
|-----|---|--------|
| N/N | 9 | 100.0% |
| M/N | 0 | 0.0% |
| M/M | 0 | 0.0% |

[Spinal Dysraphism, Weimaraner](#)

(9 tests)

| | | |
|-----|---|-------|
| N/N | 7 | 77.8% |
| M/N | 2 | 22.2% |
| M/M | 0 | 0.0% |

[MyBreedData \(Donner, 2018\)](#)

Hyperuricosuria

(72 tests)

| | | |
|-----|----|-------|
| N/N | 51 | 70.8% |
| M/N | 17 | 23.6% |
| M/M | 4 | 5.6% |

Hypomyelination

(68 tests)

| | | |
|-----|----|-------|
| N/N | 67 | 98.5% |
| M/N | 1 | 1.5% |
| M/M | 0 | 0.0% |

[Cone-Rod Dystrophy \(cord1-PRA/crd4\)](#)

(72 tests)

| | | |
|-----|----|-------|
| N/N | 70 | 97.2% |
| M/N | 2 | 2.8% |
| M/M | 0 | 0.0% |