

American Staffordshire Terrier

[\(AKC, 85 out of 204, 2022\)](#)



LABGENVET (2021)



[MyBreedData \(Donner, 2018\)](#)

[Cone Rod Dystrophy \(Cord1-PRA/crd4\)](#)

(257 tests)

N/N	181	70.4%
M/N	66	25.7%
M/M	10	3.9%

[Hyperuricosuria](#)

(258 tests)

N/N	219	84.9%
M/N	37	14.3%
M/M	2	0.8%

[Neuronal Ceroid Lipofuscinosis 4A \(NCL4\)](#)

(type Am. Staffordshire Terrier) (257 tests)

N/N	230	89.5%
M/N	25	9.7%
M/M	2	0.8%

[Degenerative myelopathy](#)

(258 tests)

N/N	234	90.7%
M/N	22	8.5%
M/M	2	0.8%

Cone Rod Dystrophy 1 (CRD1)

(type Am. Staffordshire Terrier) (242 tests)

N/N	237	97.9%
M/N	5	2.1%
M/M	0	0.0%

Cone Rod Dystrophy 2 (CRD2)
(type Pitbull Terrier) (242 tests)

N/N	238	98.3%
M/N	3	1.3%
M/M	1	0.8%

[Primary Lens Luxation \(PLL\)](#)

(258 tests)

N/N	257	99.6%
M/N	1	0.4%
M/M	0	0.0%

[Craniomandibular Osteopathy \(CMO\)](#)

(257 tests)

N/N	256	99.6%
M/N	1	0.4%
M/M	0	0.0%